

A quantum beam splitter for atoms.

Uffe V. Poulsen* and Klaus Mølmer

Institute of Physics and Astronomy, University of Aarhus, DK-8000 Århus C

An interferometric method is proposed to controllably split an atomic condensate in two spatial components with strongly reduced population fluctuations. All steps in our proposal are in current use in cold atom laboratories, and we show with a theoretical calculation that our proposal is very robust against imperfections of the interferometer.

PACS numbers: 3.75.Fi, 39.20.+q

I. INTRODUCTION

Cold atoms are routinely applied in spectroscopy and in atomic clocks, and matter wave interferometers have been used to probe with unprecedented precision the effects caused by acceleration, gravity or interaction with atoms or electromagnetic fields [1]. The experimental resolution in these experiments is linked to the precision with which the relevant signal can be measured and to the intrinsic uncertainty of the quantity measured. For example, in Ramsey interferometry applied in atomic clocks, one tunes a radiation frequency to obtain a 50 % population of both states of the clock transition, but if the atoms are uncorrelated, in addition to the measurement errors one will inevitably observe differences scaling as $\sqrt{N}$ between the populations of the two states, where N is the total number of atoms (binomial distribution). So-called spin squeezed states have been shown to exist, for which it is possible to significantly reduce this uncertainty without going to exceedingly large numbers of atoms [2, 3]. A number of proposals has been made to produce such spin squeezed states by interaction with squeezed light [4, 5], by quantum non-demolition measurement of the atomic populations [6, 7], and by light-induced [8] or collisionally induced [9, 10, 11] interaction.

In this paper we propose a method to realize a beam splitter for a one-component condensate, causing a splitting of the condensate in two spatially separated components with a better matching of occupancies than in the binomial distribution, resulting from a normal splitting of the particles. Such splitting has indeed been proposed to occur if one adiabatically raises a potential barrier inside a single condensate, and experiments have confirmed the effects of collisional interactions on the population statistics of a condensate split by a periodic potential [12]. This dynamics is governed by the time scale for which the system is able to adiabatically follow the lowest energy state due to the collisional repulsion among atoms, and this time scale may be very long, making an experimental implementation very difficult [13].

In contrast, we propose a fast method working in four steps: (i) apply a normal splitting to create a binomial




FIG. 1: Atom-interferometer realized via Bragg scattering of the condensate.

distribution of atoms separated in space, (ii) make use of the collisional interaction in each spatial component to cause in few msec a non-linear phase evolution of the different amplitudes, (iii) reflect the atoms so they again overlap in space, and (iv) remix them on a second beam splitter, so that the resulting separate components are populated by a sub-binomial distribution.

In Sec. II, we present a theoretical analysis of the interferometer applied to a Bose-Einstein condensate. In Sec. III, we discuss the results ideally obtained with our proposal, and in Sec. IV, we analyze the effect of a mismatch in overlap in the interferometer.

II. BRAGG INTERFEROMETER

The principle of the proposal is sketched in Fig. 1., where the vertical arrows indicate Bragg diffracting laser fields with appropriate detunings and phases. We suggest to use Bragg diffraction to split and to recombine the atomic clouds because this method has already been successfully demonstrated in experiments.

Our proposal combines ideas for light squeezing in a Kerr-nonlinear interferometer [14] with ideas for spin squeezing of two-component condensates using the collisional interactions among the atoms [9, 15]. In a single condensate, the collisional interaction leads to an effective quadratic term in the Hamiltonian proportional to $N(N - 1)$, where N is the total number of atoms. That term affects only the global phase of the con-

*Electronic address: uvp@ifu.au.dk

condensate, but if the condensate is divided in two components with no mutual interaction, it is replaced by $n_1(n_1 - 1) + n_2(n_2 - 1)$, having a significant effect on a superposition of states with different populations n_1 and n_2 in the two components. The lack of interaction between the components could be due to a tuning of the collisional properties of the atoms, or, as suggested in [15], due to a spatial separation of the atoms. In [15] we carried out simulations with the so-called positive-P method which retains the full many-body character of the problem. Our calculations were well accounted for by a simple two-mode description, and we shall therefore apply such a description in the following analysis.

A. Internal and external dynamics

The atoms are initially in a single component BEC, *i.e.*, all atoms populate the same one-particle wave function $\psi(\vec{r})$. Counterpropagating laser beams along the z -direction with a frequency difference around 100 kHz are applied to the atomic cloud and cause diffraction of the atoms. In the moving frame of the optical standing wave pattern, Bragg diffraction conserves kinetic energy and the atoms coherently populate two components at the incident momentum $-\hbar k$ (corresponding to zero momentum in the laboratory frame) and at $\hbar k$, differing by twice the photon momentum. For suitably chosen parameters the diffraction process is fast and interactions can be ignored. The diffraction is then a linear process and each atom is put in a superposition of remaining at rest, and having received twice the photon momentum. Allowed to propagate freely the atoms will coherently populate two spatially separated regions of space after less than a *msec*. If we neglect the role of interactions in this short phase, we can write the state of the system

$$|\Psi\rangle = \frac{1}{\sqrt{2^N}} \sum_{n_1=0}^N \sqrt{\binom{N}{n_1}} |n_1; \phi_1(\vec{r})\rangle |n_2; \phi_2(\vec{r})\rangle \quad (1)$$

where $|n_i; \phi_i(\vec{r})\rangle$ denotes a state with n_i atoms populating the spatial wave function $\phi_i(\vec{r})$; $n_2 = N - n_1$. In the subsequent dynamics, the wave functions $\phi_i(\vec{r})$ evolve with time, and we describe this evolution with the Gross-Pitaevskii equations

$$i\hbar\partial_t\phi_i = (\hat{h}_i + \frac{Ng}{2}|\phi_i|^2)\phi_i \quad (2)$$

where $\hat{h}_i$ is the single particle hamiltonian (the atoms may be free, in which case $\hat{h}_i$ equals the kinetic energy operator, they may fall under gravity, or they may be trapped in a weak trapping potential) and $g = 4\pi\hbar^2 a_s/m$ is the collisional interaction strength (a_s is the s -wave scattering length).

To compute the effect of collisions on the spatial dynamics it is an adequate approximation to assume that half of the atoms $N/2$ populate each component. A more

precise account of the interactions will be needed, however, when we examine the details of the evolution across the binomial variation in n_i . Effectively, the functions ϕ_1 and ϕ_2 define two modes for the atoms with creation and annihilation operators $a_i^\dagger$ and a_i , and the dynamics associated with the distribution of atoms among the modes is accounted for by a two-mode Hamiltonian:

$$H = \sum_{i=1,2} g\mathcal{I}_i \left(\frac{1}{2}(a_i^\dagger)^2 a_i^2 - \frac{N}{2} a_i^\dagger a_i \right) \quad (3)$$

where $\mathcal{I}_i = \int d^3r |\phi_i|^4$, is determined from the solution of Eq.(2). It is not strictly correct to separate the spatial and the population dynamics, especially not if the condensates are strongly interacting, which suggests a variation of the preferred mode function ϕ_i over the binomial distribution of n_i around $N/2$. In the Thomas-Fermi regime, this variation has been taken into account within a broader ansatz for the state of a similar two-component system suggesting, however, that this effect is taken well care of if $\mathcal{I}_i$ is multiplied by a numerical factor of order unity [16].

B. Collective spin picture

The Hamiltonian (3) will cause a quadratic (with n_i) evolution of the phase of the expansion coefficients in the state (1). This evolution is conveniently described by defining the collective spin operators

$$S_+ = a_2^\dagger a_1, \quad S_- = a_1^\dagger a_2, \quad S_z = (a_2^\dagger a_2 - a_1^\dagger a_1)/2. \quad (4)$$

The Hamiltonian (3) can now be written:

$$H = \hbar\chi S_z^2 + f(N) \quad (5)$$

with $\hbar\chi \equiv g\mathcal{I}$, where we have assumed for simplicity $\mathcal{I}_1 = \mathcal{I}_2 \equiv \mathcal{I}$. The $f(N)$ term gives an overall phase to the system which we can neglect while the quadratic term S_z^2 has been studied in detail by Kitagawa and Ueda [17], who pointed out that precisely this Hamiltonian leads to squeezing, *i.e.*, reduction of the variance of a spin component orthogonal to the direction of the mean spin as illustrated in Fig. 2.

Starting with the state (1), a spin coherent state with the macroscopic spin pointing along the x -axis, the Hamiltonian (5) produces a state which retains a large value of $\langle S_x \rangle$ and vanishing $\langle S_{y,z} \rangle$. The initial variance $\text{Var}(S_y) = N/4$ changes due to the interaction, and one identifies [17] a particular component $S_\theta = \cos\theta S_z - \sin\theta S_y$ for which the variance is minimal. In terms of $A = 1 - \cos^{N-1}(2\chi t)$ and $B = 4\sin(\chi t)\cos^{N-2}(\chi t)$ the angle θ is given by

$$\theta = \frac{1}{2} \arctan \frac{B}{A} \quad (6)$$

and the minimal variance is given by

$$\text{Var}(S_\theta) = \frac{N}{4} \left\{ 1 - \frac{N-1}{4} \left[\sqrt{A^2 + B^2} - A \right] \right\}. \quad (7)$$




FIG. 2: Phase- and population dynamics in the collective spin picture. The z -component S_z represents the population difference between the two spatial components, and the initial single component condensate starts out on the south pole with equal uncertainty in the S_x and S_y spin components. The preparation of the state with all atoms in a given superposition of the spatial modes 1 and 2 is represented by a simple rotation of the spin vector. The first Bragg pulse is a $\pi/2$ pulse and the spin is rotated to the equator of the Bloch sphere. By convention we choose the rotation axis to be the y -axis so the mean spin ends up along the x -axis. The S_z^2 term in Eq. (5) now deforms the uncertainty ellipse while leaving the mean spin along x . This deformation stretches the uncertainty ellipse in the horizontal direction while $\text{Var}(S_z)$ is kept constant. To obtain reduced fluctuations in the populations of the two spatial modes, i.e. in S_z , a rotation by the angle θ shown in the figure is applied by a final Bragg pulse, chosen with appropriate phases so that the rotation is around the direction of the mean spin.

This spin squeezing is equivalent to a collapse of the relative phase, smeared out by the interaction [16, 18, 19]. It is important to notice that the phase collapse is not a decoherent process; the system is still in a pure state, and by a simple unitary operation described below (a spin rotation, cf. Fig. (2)) it can be transformed to a state with the desired reduced population fluctuations. We focus on population fluctuations, but it is worthwhile mentioning that it is equally possible to obtain a state where the relative phase of the two components has become well defined.

The Hamiltonian (5) applies whenever the two components are spatially well separated, and we let the atomic clouds separate for a time interval of duration T (typically some *msecs*). We then apply a new Bragg diffraction pulse to induce a complete transfer between states

with momenta $\pm\hbar k$ in the moving frame, *i.e.* 0 and $2\hbar k$ in the laboratory frame, so that the two components now approach each other. This operation is carried out on the two spatially separated components, and it is accounted for by inclusion of the interaction with the Bragg fields in $\hat{h}_i$ in Eq.(2). Note that with the convention that the collective spin refers to the spatially separated components, this Bragg pulse has no effect on the values of the spin components; the solutions $\phi_i(\vec{r}, t)$ constitute a 'rotating frame' for our calculation of populations and coherences.

After a second time interval of duration T with free evolution of the two condensate components, they again overlap in space. At this point it is possible to observe interference fringes in the spatial density profile [20, 21]. Note that although Fig.2 is suggestive that spin measurements and rotations can be carried out at any time, it is only when the two spatial components overlap, the spin variables represent physically accessible quantities. At the time of full overlap we can apply a final beam splitter Bragg pulse to recombine the two components into two new output components:

$$U_{\text{Bragg}} = \exp[i\alpha(\cos\phi S_x + \sin\phi S_y)] \\ = \begin{bmatrix} \cos\frac{\alpha}{2} & i\sin\frac{\alpha}{2}e^{-i\phi} \\ i\sin\frac{\alpha}{2}e^{i\phi} & \cos\frac{\alpha}{2} \end{bmatrix} \quad (8)$$

Here we have written the Bragg evolution operator in the spin picture and as a 2×2 matrix giving the mode annihilation operators b_1 and b_2 after the pulse in terms of the operators a_1 and a_2 before the pulse. We are interested in creating a state with two separate components with sub-binomial counting statistics, that is, we want to choose ϕ , the phase of the pulse, and α , the duration, such that the counting statistics of b_1 and b_2 will benefit from the correlations in a_1 and a_2 . In the spin picture it is clear that this operation should be a rotation around the mean spin to align the squeezed direction with the z -axis which represents population differences. If we denote by S_α the spin component that the last Bragg pulse rotates into the z -direction then from the definition (4) we get the mean and variance of the populations of one of the output beams

$$\langle n_{\text{det}} \rangle = \frac{1}{2}N + \langle S_\alpha \rangle \\ \text{Var}(n_{\text{det}}) = \text{Var}(S_\alpha). \quad (9)$$

If $\phi = 0$ we rotate around the x -axis and the angle α should ideally be chosen equal to θ of Fig. 2 and Eq. (6). With no spin squeezing one finds $\text{Var}(n_{\text{det}}) = N/4$ in agreement with the initial binomial distribution.

From Eq. (8) it is clear that the distinction between different rotation axes, all lying in the xy -plane is equivalent to the one between complex phases ϕ of the coupling amplitude, *i.e.* the relative phase of the two counterpropagating laser beams creating the moving standing wave. Alternatively, we see that the location of the Bragg diffraction pattern, due to the translational properties of momentum eigenstates, controls precisely the

phase of the coupling. In Ref. [15], we learned that the mean spin and the rotation angle are predicted with adequate precision in the simple model (5,6), so that application of Bragg pulses with these variables should suffice to yield substantial reduction of the population fluctuations of the two atomic outputs. It is also straightforward to optimize the parameters in experiments. The phases of the diffracting lasers are adjusted so that the two output beams have the same mean occupancy, independently of the duration of the coupling. The solid curve in Fig. 3 shows the squeezing factor $(N/4)/\text{Var}(n_{\text{det}})$ (large for strong squeezing) as a function of α . With a spin squeezed sample the variance of the populations show a strong dependence on the duration of the last Bragg pulse, which should therefore be adjusted to identify the output with minimal fluctuations.

III. EXPECTED RESULTS

Let us turn to the discussion of the experimental feasibility of our proposal. Splitting and recombination of one-component condensates have been done in several laboratories, and the coherence properties have been verified and explicitly utilized in a number of imaging experiments: our proposal follows closely the experiments at NIST, where the phase variation of the condensate gives rise to a density variation in the recombined output [22]. These experiments have been done in the limit of interacting condensates which is of course essential for our proposal: In [21] the mean field repulsion was observed and it was shown not to prevent a nearly perfect overlap of the spatial modes at the recombination, and in [22] a soliton was imprinted in one of the components to be subsequently detected in the output. We note, that also the phase variation around quantized vortices has been studied by similar interference imaging, both in the case where a vortex was prepared prior to the splitting and a dislocation appears in the interference fringes [23], and in the case where a condensate was split and a vortex was subsequently created by stirring only one component [24].

To assess the strength of the squeezing interaction we need to evaluate the parameter χ of Eq.(5). In the Thomas-Fermi approximation, valid for an interaction dominated condensate, one finds the simple result:

$$\mathcal{I} = \int |\phi|^4 d^3r = \frac{10}{7} \frac{1}{V_{\text{TF}}} \quad (10)$$

where $V_{\text{TF}} \equiv (4\pi/3)R_x R_y R_z$ with R_i the Thomas-Fermi radius of the condensate in the i -direction. With the parameters of the experiments with condensates, being split and recombined by Bragg diffraction at NIST, [22], one finds $\chi = 4.7 \times 10^{-4} \text{s}^{-1}$ or a factor of ~ 4 reduction in the population variances in the 2 *msec* duration of the experiment. This result is improved to a factor of more than 12 by increasing the duration to 4 *msec*. The theoretical maximal squeezing factor

$\sim (1/3)(N/6)^{-2/3} \sim 1.3 \times 10^5$ [17] would require the components to be separated for $\sim 6^{1/6} \chi N^{-2/3} \sim 200 \text{msec}$ assuming a constant χ throughout the experiment. In reality the condensate components expand if the trap is turned off during the experiment and the continuous reduction of χ over longer times according to Eq.(10) would have to be determined by an integration of the Gross-Pitaevskii equation (2).

IV. EFFECT OF IMPERFECT OVERLAP

As a final issue we wish to discuss the matching of the spatial wave functions of the two components. After the physical interactions described above, one will detect a number of atoms with momenta around zero, and a number of atoms with momenta around $2\hbar k$, and the analysis shows that these numbers will fluctuate less than if they were given by a binomial distribution. This reduction is due to the correlations between the atoms following the lower and the upper paths in the interferometer, and it is hence important that atoms from the lower path, being diffracted by the last Bragg pulse, occupy the same spatial state as the undiffracted component of the upper path, cf. Fig. 1. In the extreme case where these two contributions are orthogonal and distinguishable, there will be no squeezing effect at all, and in the case where they overlap 100 %, the squeezing is given by the above simple analysis.

Atoms leaving the interferometer at zero momentum can in general be described by a superposition of ϕ_2 , the undiffracted upper path wave function, and a state $\phi_{2\perp}$ orthogonal to ϕ_2 . This gives rise to two orthogonal modes which both contribute to the detected number of atoms around zero momentum. We can express the field annihilation and creation operators $b_2, b_{2\perp}$ and $b_2^\dagger, b_{2\perp}^\dagger$ for these two modes in terms of the operators a_1, a_2 and $a_1^\dagger, a_2^\dagger$ given above for the modes prior to the last Bragg pulse *and* operators for two additional 'vacuum' modes $a_{1\perp}, a_{2\perp}$ and $a_{1\perp}^\dagger, a_{2\perp}^\dagger$, which are necessary to ensure unitarity of the beam splitter pulse. Letting c denote the overlap of the (normalized) diffracted lower path component with the upper path undiffracted component and s the overlap with the orthogonal complement

$$e^{-2ikx} \phi_1(\vec{r}) = c\phi_2(\vec{r}) + s\phi_{2\perp}(\vec{r}) \quad (11)$$

($|s|^2 + |c|^2 = 1$). In terms of α and ϕ of the last Bragg pulse (8) we have:

$$b_2 = \cos \frac{\alpha}{2} a_2 + i \sin \frac{\alpha}{2} e^{-i\phi} (c a_1 + s a_{1\perp}) \quad (12)$$

$$b_{2\perp} = \cos \frac{\alpha}{2} a_{2\perp} + i \sin \frac{\alpha}{2} e^{-i\phi} (s^* a_1 - c^* a_{1\perp}). \quad (13)$$

Focusing on b_2 we see that for $|c| < 1$ we can still control the mixing and pick the squeezed combination of a_1 and a_2 . However, some contributions from the unoccupied modes have been introduced, and the situation is similar

to the detection of squeezed light: A non-unit overlap ν between the squeezed mode of light and a detector mode as, e.g., modelled by a beam splitter transmitting a fraction $|\nu|^2$ of the squeezed light to the detector, leads to admixture with a vacuum field and the deterioration of the squeezing, $\text{Var}(X)_{\text{det}} = |\nu|^2 \text{Var}(X)_{sq} + (1 - |\nu|^2)$, where the X is the scaled quadrature field component with unit variance in the vacuum state.

In a real experiment it is difficult to count the atoms in mode ϕ_2 exclusively. It is more realistic that one will only have access to the total number of atoms in ϕ_2 and $\phi_{2\perp}$. We then have to include $b_{2\perp}$ in the analysis, and if the phase of the Bragg diffraction is chosen to compensate the phase of the complex overlap c , we find:

$$\begin{aligned} \langle n_{\text{det}} \rangle &= \langle b_2^\dagger b_2 + b_{2\perp}^\dagger b_{2\perp} \rangle \\ &= \langle \cos \alpha S_z - |c| \sin \alpha S_y \rangle + \frac{N}{2} \\ &= \lambda \langle S_{\alpha'} \rangle + \frac{N}{2}. \end{aligned} \quad (14)$$

in terms of the spin component $S_{\alpha'}$ prior to the last Bragg pulse. The angle α' is related to α , the rotation angle given by the duration of the last Bragg pulse, by $\tan \alpha' = |c| \tan \alpha$, and we have introduced the factor $\lambda = \sqrt{|c|^2 / (\sin^2 \alpha' + |c|^2 \cos^2 \alpha')}$.

The spin component $S_{\alpha'} \equiv \cos \alpha' S_z - \sin \alpha' S_y$ is in a direction perpendicular to the mean spin. We therefore have $\langle S_{\alpha'} \rangle = 0$ and $\langle n_{\text{det}} \rangle = N/2$ independently of α . The variance of n_{det} is given by

$$\begin{aligned} \text{Var}(n_{\text{det}}) &= \langle n_{\text{det}}^2 \rangle - \langle n_{\text{det}} \rangle^2 \\ &= \lambda^2 \text{Var}(S_{\alpha'}) + (1 - \lambda^2) \frac{N}{4}. \end{aligned} \quad (15)$$

If the overlap is complete $|c| = 1 = \lambda$, we can pick out the maximally squeezed spin component to yield the minimal variance of n_{det} . In case of a non-perfect overlap, we see that Eq. (15) introduces a non-squeezed contribution, scaling as $N/4$, and it makes the number fluctuations depend on the noise of the spin component $S_{\alpha'}$ rather than of S_α . We therefore want α' to be close to the actual direction of squeezing θ in Fig.2 and Eq.(6), which is obtained by choosing a Bragg pulse with longer duration than what is optimal in the ideal case.

In Fig. 3 we show the variation of $(N/4)/\text{Var}(n_{\text{det}})$ with α for different values of the overlap c . When $|c|$ is reduced the squeezing factor drops, but the curves illustrate that the results improves if one choses a larger rotation angle α .

Note that the mode function mismatch does not have the same detrimental effect as in the detection of a single mode squeezed field: the admixture of the vacuum modes, and hence the vacuum contribution $N/4$ to $\text{Var}(n_{\text{det}})$, does not scale with $|s|^2 = 1 - |c|^2$ but rather with $|s|^2 \sin^2 \alpha$, and for small rotation angles this is a small number. For strong squeezing the uncertainty ellipse is almost horisontal and only rotation by a small




FIG. 3: The reduction of $\text{Var}(n_{\text{det}})$ is illustrated by plotting the reciprocal ratio $(N/4)/\text{Var}(n_{\text{det}})$ as a function of the applied Bragg pulse rotation angle for parameters as in Sec. III with 4 msec separation. The results are shown for different values of the overlap, $c = 1$ (solid line), $c = 0.9$ (dashed line), $c = 0.8$ (dash-dotted curve), and $c = 0.6$ (dotted curve). The value of α at the maximum of the $c = 1$ curve displays the orientation of the squeezing ellipse, i.e. θ of Fig. 2. As is apparent from the other curves, one may compensate for a non-perfect overlap by applying a larger Bragg pulse rotation.

α is needed to observe squeezing in the number fluctuations. In this case even a significantly reduced overlap does not prevent a significant noise reduction for n_{det} . This is illustrated when we write the maximal squeezing factor attainable for given c and given $\text{Var}(S_\theta)$:

$$\begin{aligned} \text{Var}(n_{\text{det}}) &= \text{Var}(S_\theta) \left(1 + (1 - |c|^2) \frac{\frac{N}{4} - \text{Var}(S_\theta)}{|c|^2 \frac{N}{4} + \text{Var}(S_\theta)} \right) \\ &\sim \frac{\text{Var}(S_\theta)}{|c|^2} \quad (\text{Var}(S_\theta) \ll \frac{N}{4}). \end{aligned} \quad (16)$$

(In Eq. (16) we assume that the spin state described by Eqs. (6,7) is a minimum uncertainty state).

V. CONCLUSION

We have proposed to use the collisional interactions between atoms in the separate arms of a spatial interferometer to suitably manipulate the amplitudes on the different number state components. We analyzed the achievements of current condensate interferometry experiments, and we found that significant number squeezing should be found in the interferometer output, i.e., the interferometer acts as a quantum beam splitter for atoms with unique noise properties. Good overlap of the output mode functions is necessary to observe interference

effects in the interferometer and it is also important for the noise reduction in our beam splitter. We showed, however, that the overlap is not a critical parameter and

sizable noise reduction is possible with interferometric overlaps easily achievable in experiments.

-
- [1] P. Berman, ed., *Atom Interferometry* (Academic Press, New York, 1997).
 - [2] D. J. Wineland, J. J. Bollinger, W. M. Itano, and D. J. Heinzen, *Phys. Rev. A* **50**, 67 (1994).
 - [3] G. S. Agarwal and R. R. Puri, *Phys. Rev. A* **41**, 3782 (1990).
 - [4] A. Kuzmich, K. Mølmer, and E. S. Polzik, *Phys. Rev. Lett.* **79**, 4782 (1997).
 - [5] J. Hald, J. L. Sørensen, C. Schori, and E. S. Polzik, *Phys. Rev. Lett.* **83**, 1319 (1999).
 - [6] A. Kuzmich, L. Mandel, and N. P. Bigelow, *Phys. Rev. Lett.* **85**, 1594 (2000).
 - [7] K. Mølmer, *Euro. Phys. J. D* **5**, 301 (1999).
 - [8] I. Bouchoule and K. Mølmer (2001), submitted for publication in *Phys. Rev. Lett.*, quant-ph/0105144.
 - [9] A. Sørensen, L.-M. Duan, I. Cirac, and P. Zoller, *Nature* **409**, 63 (2001).
 - [10] J. A. Dunningham, K. Burnett, and M. Edwards, *Phys. Rev. A* **64**, 015601 (2001).
 - [11] S. Raghavan, H. Pu, P. Meystre, and N. Bigelow, *Opt. Comm.* **188**, 149 (2001).
 - [12] C. Orzel, A. K. Tuchman, M. L. Fenselau, M. Yasuda, and M. A. Kasevich, *Science* **291**, 2386 (2001).
 - [13] C. Menotti, J. R. Anglin, J. I. Cirac, and P. Zoller, *Phys. Rev. A* **63**, 023601 (2001).
 - [14] M. Kitagawa and Y. Yamamoto, *Phys. Rev. A* **34**, 3974 (1986).
 - [15] U. V. Poulsen and K. Mølmer, *Phys. Rev. A* **64**, 013616 (2001), quant-ph/0101089.
 - [16] A. Sinatra and Y. Castin, *Euro. Phys. J. D* **8**, 319 (2000).
 - [17] M. Kitagawa and M. Ueda, *Phys. Rev. A* **47**, 5138 (1993).
 - [18] J. Javanainen and M. Wilkens, *Phys. Rev. Lett.* **78**, 4675 (1997).
 - [19] K. Mølmer, *Phys. Rev. A* **58**, 566 (1998).
 - [20] M. R. Andrews, C. G. Townsend, H.-J. Miesner, D. S. Durfee, D. M. Kurn, and W. Ketterle, *Science* **275**, 637 (1997).
 - [21] J. E. Simsarian, J. Denschlag, M. Edwards, C. W. Clark, L. Deng, E. W. Hagley, K. Helmerson, S. L. Rolston, and W. D. Phillips, *Phys. Rev. Lett.* **85**, 2040 (2000), cond-mat/0005303.
 - [22] J. Denschlag, J. E. Simsarian, D. L. Feder, C. W. Clark, L. A. Collins, J. Cubizolles, L. Deng, E. W. Hagley, K. Helmerson, W. P. Reinhardt, et al., *Science* **287**, 97 (2000).
 - [23] F. Chevy, K. W. Madison, V. Bretin, and J. Dalibard (2001), cond-mat/0104545.
 - [24] S. Inouye, S. Gupta, T. Rosenband, A. Chikkatur, A. Görlitz, T. Gustavson, A. Leanhardt, D. Pritchard, and W. Ketterle (2001), cond-mat/0104444.